

Complete Genome Sequence of the Rearranged Porcine Circovirus Type 2

Libin Wen, Kongwang He, Yanxiu Ni, Xuehan Zhang, Bin Li,
Xiaomin Wang, Rong li Guo, Zhengyu Yu, Aihua Mao,
Junming Zhou, Lixin Lv and Jieyuan Jiang
J. Virol. 2012, 86(10):5963. DOI: 10.1128/JVI.00494-12.

Updated information and services can be found at:
<http://jvi.asm.org/content/86/10/5963>

	<i>These include:</i>
REFERENCES	This article cites 5 articles, 1 of which can be accessed free at: http://jvi.asm.org/content/86/10/5963#ref-list-1
CONTENT ALERTS	Receive: RSS Feeds, eTOCs, free email alerts (when new articles cite this article), more»

Information about commercial reprint orders: <http://jvi.asm.org/site/misc/reprints.xhtml>
To subscribe to to another ASM Journal go to: <http://journals.asm.org/site/subscriptions/>

Complete Genome Sequence of the Rearranged Porcine Circovirus Type 2

Libin Wen, Kongwang He, Yanxiu Ni, Xuehan Zhang, Bin Li, Xiaomin Wang, Rong li Guo, Zhengyu Yu, Aihua Mao, Junming Zhou, Lixin Lv, and Jieyuan Jiang

Institute of Veterinary Medicine, Jiangsu Academy of Agricultural Sciences, Key Laboratory of Veterinary Biological Engineering and Technology, Ministry of Agriculture, National Center for Engineering Research of Veterinary Bio-products, Nanjing, China

We first report here the genome sequences of 4 rearranged porcine circovirus type 2 strains, JSTZ, ZJQDH1, ZJQDH2, and JSHM, isolated from porcine sera in China. The complete circular genomes of these isolates are 578, 483, 574, and 772 nucleotides in length, respectively. They are predicted to be defective interfering particles of porcine circovirus type 2. The findings will help us to understand molecular evolution of porcine circovirus type 2 and the relationship between porcine circovirus type 2 and diseases.

Porcine circovirus (PCV), a small, nonenveloped, single-stranded DNA virus with a circular genome from the *Circoviridae* family, was first described as a picornavirus-like contaminant of the continuous porcine kidney cell line PK-15 (ATCC CCL31) in 1974 (3).

The PK-15-derived PCV isolate, designated PCV1, has 1,759 nucleotides (nt) and is considered nonpathogenic (2), while a novel PCV isolate, namely, PCV2, with 1,768 or 1,767 nt, has been considered the primary etiological agent, although not the sole factor, in developing a group of complex multifactorial PCV diseases (PCVD) or PCV-associated diseases (PCVAD), especially postweaning multisystemic wasting syndrome (PMWS). PMWS was first recognized in North America in 1991 (1). Since then, it has caused severe losses in pig production worldwide.

We isolated and identified 4 novel rearranged PCV2 isolates from pigs suffering from PMWS in China. All 4 genomic sequences analyzed here were obtained by PCR analysis and sequenced with an ABI 3730 genome sequencer. Overlapping consensus sequences were assembled and open reading frames (ORFs) were identified using DNAMAN version 5.2.2.

The complete circular genomes of the four rearranged PCV2 isolates, JSTZ, ZJQDH1, ZJQDH2, and JSHM, are 578, 483, 574, and 772 nt in length, respectively. These DNA molecules are sub-genomic rearranged molecules originating from the parental PCV2 genome and display similar structures: they contain the origin of PCV2 genome replication, characterized by a putative stem-loop structure, with a conserved nonanucleotide motif (AA GTATTAC) and 2 to 3 hexamer motifs (CGGCAG) serving as binding sites for the replicases. They include the predicted ORF encoding a whole or truncated capsid protein of PCV2.

Unlike the porcine-circovirus-like agents P1 and P2 (4, 5, 6), the four rearranged PCV2 genomes do not contain the insertion of the exogenous sequence, so they are predicted to be defective interfering particles of PCV2. Of course, it remains to be clarified

whether these rearranged components resulting from PCV2 may contribute to PMWS.

Nucleotide sequence accession numbers. The complete genome sequences of the four rearranged PCV2 isolates, JSTZ, ZJQDH1, ZJQDH2, and JSHM, have been deposited in GenBank under the accession numbers [JQ690758](#), [JQ690759](#), [JQ690760](#), and [JQ690761](#), respectively.

ACKNOWLEDGMENTS

This work was supported by grants from the National Natural Science Foundation of China (30972184) and the Jiangsu Provincial Natural Science Foundation (BK2008351).

REFERENCES

1. Clark EG. 1997. Post-weaning multisystemic wasting syndrome, p 499–501. *In* Proceedings of the 28th Annual Meeting of the American Association of Swine Practitioners. American Association of Swine Practitioners, Quebec, Canada.
2. Tischer I, Miels W, Wolff D, Vagt M, Griem W. 1986. Studies on epidemiology and pathogenicity of porcine circovirus. *Arch. Virol.* 91:271–276.
3. Tischer I, Rasch R, Tochtermann G. 1974. Characterization of papovavirus- and picornavirus-like particles in permanent pig kidney cell lines. *Zentralb. Bakteriol.* 226:153–167.
4. Wen L, et al. 2008. Complete nucleotide sequence of a novel porcine circovirus-like agent and its infectivity in vitro. *Sci. China C Life. Sci.* 51(5): 453–458.
5. Wen L, et al. 2011. Prevalence of porcine circovirus-like agent P1 in Jiangsu, China. *Virol. J.* 8:543.
6. Wen L, et al. 2012. Complete genome sequence of a novel porcine circovirus-like agent. *J. Virol.* 86:639.

Received 27 February 2012 Accepted 29 February 2012

Address correspondence to Kongwang He, kwh2003@yeah.net.

Copyright © 2012, American Society for Microbiology. All Rights Reserved.

[doi:10.1128/JVI.00494-12](https://doi.org/10.1128/JVI.00494-12)